

[Letter to the Editor] 17q21 variant increases the risk of exacerbations in asthmatic children despite inhaled corticosteroids use

Article (Published Version)

Farzan, N, Vijverberg, S J, Hernandez-Pacheco, N, Bel, E H D, Berce, V, Bønnelykke, K, Bisgaard, H, Burchard, E G, Canino, G, Celedón, J C, Chew, F T, Chiang, W C, Cloutier, M M, Forno, E, Francis, B et al. (2018) [Letter to the Editor] 17q21 variant increases the risk of exacerbations in asthmatic children despite inhaled corticosteroids use. *Allergy*, 73 (10). pp. 2083-2088. ISSN 0105-4538

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/76437/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

17q21 variant increases the risk of exacerbations in asthmatic children despite inhaled corticosteroids use

To the Editor,

Approximately 25% of the asthmatic children suffer from uncontrolled asthma despite regular use of inhaled corticosteroids (ICS).¹ Variation within the 17q21 locus is the strongest genetic determinant for childhood-onset asthma.² Recently, the influence of this locus on treatment outcomes has been shown in several studies.³⁻⁴ The Pharmacogenomics in Childhood Asthma (PiCA) consortium is a multiethnic consortium that brings together data from $\geq 14\,000$ asthmatic children/young adults from 12 different countries to study the pharmacogenomics of uncontrolled asthma despite treatment.⁵ In 14 PiCA populations (with over 4000 asthmatic patients), we studied the association between variation in the 17q21 locus, and asthma exacerbations despite ICS use. We specifically focused on rs7216389, a single nucleotide polymorphism (SNP) in the 17q21 locus strongly associated with childhood asthma and initially identified by Moffatt et al.²

Ten PiCA studies included patients with non-Hispanic European origins, two included Hispanic patients, one African American, and one included East Asian patients. Additional details of the study populations can be found in the Data S1. Two outcomes were assessed: (i) asthma-related hospitalizations/emergency department visit (ED) visits and (ii) short courses of oral corticosteroid (OCS) use reported by the parent/child at the study visit or based on completed study questionnaires. Age, gender, genotype data, and exacerbation data were available for 4529 steroid-treated children and young adults (Table 1). Logistic regression analysis was used to assess the risk of exacerbations when carrying rs7216389. Due to potential heterogeneity between cohorts, the odds ratios (ORs) were meta-analyzed with the inverse variance weighting method assuming random effects. See Data S1 for more detail.

The risk allele (T) frequency was highest in East Asians ($n = 182$, $T = 0.81$), followed with African Americans ($T = 0.79$, $n = 468$) and Hispanics ($T = 0.66$, total $n = 916$), and it was less frequent in patients with European ancestry (ranged between 0.54 and 0.62, total $n = 2963$). The genotype distribution of the SNP was in Hardy-Weinberg equilibrium in all cohorts. There was a low to moderate heterogeneity between studies (Figure 1). Exacerbation rates ranged between 6.5% (PACMAN) and 77.2% (HPR) for OCS use and 6% (PACMAN) and 58% (GALA II and HPR) for hospitalizations/ED visits.

Thirty percent (1378 out of 4454) of the patients reported hospitalizations/ED visits. In the meta-analysis of 13 studies, rs7216389 was statistically significantly associated with asthma-related ED visits/hospitalizations, (summary OR per increase in risk allele: 1.32, 95% CI: 1.17- 1.49, $P < .0001$, $I^2 = 3.9\%$) (Figure 1A). In a subgroup analysis, the effect estimates for hospitalizations/ED visits were approximately the same for both non-Hispanic whites ($n = 2888$, OR: 1.33, 95% CI: 1.10-1.61, $P = .004$, $I^2 = 30.2\%$) and Hispanics ($n = 916$, OR: 1.31, 95% CI: 1.06-1.63, $P = .01$, $I^2 = 0.00\%$). Thirty-one percent (1269 out of 4050) of the patients reported OCS use/high-dose ICS. In the meta-analysis of the nine studies, the rs7216389-T was statistically significantly associated with an increased risk of OCS use/high-dose ICS (summary OR per increase in variant allele: 1.19, 95% CI: 1.04-1.36, $P = .01$, $I^2 = 22.8\%$) (Figure 1B). Rs7216389 was associated with OCS use in the meta-analysis of seven European studies ($n = 2492$, OR: 1.26, 95% CI: 1.09-1.45, $P = .002$, $I^2 = 6.2\%$) but not in Hispanics ($n = 916$, OR: 0.96, 95% CI: 0.76-1.22, $P = .7$, $I^2 = 0.00\%$). Differences in the minor allele frequencies and LD structures among different ethnicities can influence results of the association studies.⁶ This could be one of the potential explanations why we did not find a significant association in African Americans and patients from Singapore.

A sensitivity analysis was performed to investigate this association in children ≥ 5 years of age. When excluding children < 5 years of age in the meta-analysis, the results remained significant. In the meta-analysis of 13 studies, the SNP was associated with asthma-related hospitalization/ED visits ($n = 4254$, OR: 1.32, 95% CI: 1.18-1.49, $P < .0001$, $I^2 = 0.0\%$) (Figure S1). Regarding OCS use, 10 studies collected data on patients ≥ 5 years of age ($n = 3771$). In the meta-analysis of 10 studies, rs7216389 was associated with the OCS use (summary OR: 1.20, 95% CI: 1.05-1.38, $P = .01$, $I^2 = 21.7\%$) (Figure S2).

We also performed a meta-analysis of the studies that had sufficient data available on preschool children (2-4 years of age). Although the effect estimates in younger children were in the same direction for both outcomes, the results were not statistically significant (Figures S3 & S4). All preschool studies solely included European children.

Altered expression of *ORMDL3* and *GSDMB* by 17q21 locus variants may play a key role in childhood asthma onset.^{2,7} Two 17q21 asthma-risk variants (rs4065275 and rs12936231) in high

Farzan and Vijverberg equally contributed to this study.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2018 The Authors. *Allergy* Published by John Wiley & Sons Ltd.


TABLE 1 Characteristics of the study populations

	BAMSE ^a (n = 122)	BREATHE (n = 808)	CAMP ^a (n = 172)	COPSAC2000 ^{a,b} (n = 54)	ESTaTe (n = 95)	followMAGICS ^a (n = 150)	GALA II ^a (n = 744)	HPR ^a (n = 172)	PACMAN (n = 665)	PAGES (n = 308)	PASS ^c (n = 390)	SAGE II ^a (n = 468)	SCSGES ^a (182)	Slovenia ^a (n = 199)
Patient characteristic														
Age (y), mean (SD)	8.37 (0.41)	9.8 (4.0)	8.8 (2.1)	3.3 (1.0)	10.8 (4.3)	17.13 (3.03)	12.1 (3.2)	9.8 (2.7)	8.7 (2.3)	9.2 (3.8)	11.1 (4.0)	13.5 (3.4)	13.35 (5.09)	10.9 (3.41)
Male gender, % (n)	79 (96)	60.8 (491)	55.2 (95)	54 (29)	57.9 (55)	59.3 (89)	56.8 (423)	49.4 (85)	61.1 (406)	56.8 (175)	55.9 (218)	54.1 (253)	67.6 (123)	54.8 (109)
Asthma exacerbations in past year														
Asthma-related ED visit/ hospital admission, % (n) ^a	14.7 (18)	19.0 (154)	13.4 (23)	–	10.5 (10)	8.6 (13)	58.3 (434)	58.1 (100)	6.0 (39/644)	15.6 (48)	76 (296)	44.7 (209)	20.3 (37)	35.6 (71)
Oral corticosteroid use, % (n)	–	31.7 (256)	47.1 (81)	11.1 ^b (6)	36.8 (35)	–	42.3 (315)	77.3 (133)	6.5 (43)	43.2 (133)	52 (203)	29.3 (137)	20.3 (37)	–
BTS treatment step														
2, %	–	65.6	– ^c	–	60	28.7	41.1	60.7	72	25.6	–	68.6	–	–
3, %	–	18.4	–	–	37.9	60.7	43.6	36.4	22.3	61	–	25	–	–
4, %	–	16.0	–	–	2.1	10.6	15.3	2.9	5.7	13.3	–	6.4	–	–
Rs7216389														
T-allele frequency	0.57	0.56	0.58	0.62	0.60	0.58	0.66	0.66	0.59	0.58	0.60	0.79	0.81	0.54

–, data not available; BTS, British Thoracic Society; ED, emergency department; SD, standard deviation; y, years.

^aDNA was extracted from peripheral blood samples in these studies, and in the remaining, DNA was extracted from saliva samples.^bPatients with exacerbations were treated with oral corticosteroid or high-dose inhaled corticosteroids.^cCAMP is randomized clinical trial of mild to moderate asthmatics, and all children were on 200 µg of budesonide (ICS) plus SABA as needed.

(A)



(B)




FIGURE 1 Forest Plots of rs7216389 for (A) asthma-related hospitalizations/ED visits in thirteen studies and (B) OCS/high-dose ICS use in eleven studies. Odds Ratios (OR) and corresponding 95% Confidence Intervals (95% CI) for individuals with rs7216389, controlling for age, gender, and BTS treatment step

linkage disequilibrium (LD) with rs7216389 were reported to switch CTCF-binding sites that resulted in increased expression of ORMDL3 in CD4+ T cells which subsequently reduced interleukin-2 production.⁸

Rs7216389 has previously shown to be associated with exacerbations³ and poor lung function in Caucasian children using ICS.⁴ Even though in our study Caucasians were the largest group, this study is the largest multiethnic population evaluating the association between 17q21 variant and asthma exacerbations in ICS users. Rs7216389 seems to increase bronchial responsiveness and therefore exacerbation rates in children,⁹ suggesting that carriers of rs7216389 might have a more severe form of asthma. However, by adding British Thoracic Society (BTS) treatment steps as a marker of disease severity to the model, we argue that the association reflects, at least partly, poor response to ICS.

Limitations of the study include the use of retrospective reporting of exacerbations in the observational cohort studies. However, the effect was also observed in a clinical trial population (CAMP), where exacerbations were reported prospectively. Hence, we do not believe that using retrospective data has significantly influenced the results. As not all studies had data available on both hospitalizations/ED visits and OCS use, we did not combine the two outcomes in our analysis. Furthermore, as information on treatment adherence was not available in all included studies, it was not considered in the analysis.













We show that 17q21, a widely replicated asthma susceptibility locus, is also associated with an increased risk of exacerbations in children/young adults treated with ICS. A better understanding of the molecular mechanisms underlying exacerbation-prone phenotype of pediatric asthma could lead to a better classification of different pediatric asthma phenotypes and the identification of novel treatment targets.














CONFLICTS OF INTEREST

AHM reports an unrestricted research grant from GSK, during the conduct of the study; she was a member of an advisory board for AstraZeneca, outside the submitted work. MP-Y reports grants from Spanish Ministry of Economy and Competitiveness (RYC-2015-17205), from Instituto de Salud Carlos III (ISCIII, AC15/00015), and from ERA-PerMed 1st Joint Transnational Call (SysPharmPedia), during the conduct of the study. NHP reports grants from Instituto de Salud Carlos III (ISCIII) and cofunded by the European Social Funds from the European Union (ESF) "ESF invests in your future", during the conduct of the study. KGT reports grants from U.S. National Institutes of Health, during the conduct of the study. SJV reports grants from Stichting Astma bestrijding, during the conduct of the study; and PACMAN cohort was funded by a strategic alliance between Utrecht Institute for Pharmaceutical Sciences and GSK. The other authors have no other conflict of interests that are directly relevant to the content of this manuscript.

ORCID

N. Farzan  <http://orcid.org/0000-0002-3694-1086>
 S. J. Vijverberg  <http://orcid.org/0000-0002-4579-4081>
 V. Berce  <http://orcid.org/0000-0002-0577-8925>
 H. Bisgaard  <http://orcid.org/0000-0003-4131-7592>
 E. G. Burchard  <http://orcid.org/0000-0001-7475-2035>
 J. C. Celedón  <http://orcid.org/0000-0003-1366-5936>
 F. T. Chew  <http://orcid.org/0000-0003-1337-5146>
 E. Forno  <http://orcid.org/0000-0001-6497-9885>
 B. Francis  <http://orcid.org/0000-0002-2130-5976>
 D. B. Hawcutt  <http://orcid.org/0000-0002-8120-6507>
 M. Kabesch  <http://orcid.org/0000-0003-0697-1871>
 S. K. Merid  <http://orcid.org/0000-0001-5974-7676>
 M. Pino-Yanes  <http://orcid.org/0000-0003-0332-437X>
 M. Pirmohamed  <http://orcid.org/0000-0002-7534-7266>
 U. Potočník  <http://orcid.org/0000-0001-8688-174X>
 K. Repnik  <http://orcid.org/0000-0003-0801-1911>
 M. Schieck  <http://orcid.org/0000-0001-5878-0546>
 A. Sevelsted  <http://orcid.org/0000-0001-7117-2334>
 R. L. Smyth  <http://orcid.org/0000-0003-1406-6142>
 P. Soares  <http://orcid.org/0000-0001-5033-9115>
 C. Söderhäll  <http://orcid.org/0000-0002-8397-3080>
 S. M. Tse  <http://orcid.org/0000-0002-0295-0064>
 S. Turner  <http://orcid.org/0000-0001-8393-5060>
 K. M. Verhamme  <http://orcid.org/0000-0001-8162-4904>
 A.-H. Maitland-van der Zee  <http://orcid.org/0000-0002-6261-9445>

N. Farzan^{1,2} 
 S. J. Vijverberg^{1,2} 
 N. Hernandez-Pacheco^{3,4}
 E. H. D. Bel²
 V. Berce^{5,6} 
 K. Bønnelykke⁷
 H. Bisgaard⁷ 
 E. G. Burchard⁸ 
 G. Canino⁹
 J. C. Celedón¹⁰ 
 F. T. Chew^{11,12} 
 W. C. Chiang¹²
 M. M. Cloutier¹³
 E. Forno¹⁰ 
 B. Francis¹⁴ 
 D. B. Hawcutt^{14,15} 
 E. Herrera-Luis⁴
 M. Kabesch¹⁶ 
 L. Karimi¹⁷
 E. Melén^{18,19}
 S. Mukhopadhyay^{20,21}
 S. K. Merid¹⁸ 
 C. N. Palmer²¹

M. Pino-Yanes^{3,4,22} 
 M. Pirmohamed²³ 
 U. Potočnik^{5,24} 
 K. Repnik^{5,24} 
 M. Schieck^{16,25} 
 A. Sevelsted⁷ 
 Y. Y. Sio^{11,12}
 R. L. Smyth²⁶ 
 P. Soares²⁰ 
 C. Söderhäll^{27,28} 
 K. G. Tantisira^{29,30}
 R. Tavendale²¹
 S. M. Tse³¹ 
 S. Turner³² 
 K. M. Verhamme¹⁷ 
 A.-H. Maitland-van der Zee^{1,2} 

¹Division of Pharmacoepidemiology and Clinical Pharmacology, Faculty of Science, Utrecht University, Utrecht, The Netherlands

²Department of Respiratory Medicine, Academic Medical Center (AMC), University of Amsterdam, Amsterdam, The Netherlands

³Research Unit, Hospital Universitario N.S. de Candelaria, Universidad de La Laguna, Santa Cruz de Tenerife, Spain

⁴Genomics and Health Group, Department of Biochemistry, Microbiology, Cell Biology and Genetics, Universidad de La Laguna, La Laguna, Tenerife, Spain

⁵Centre for Human Molecular Genetics and Pharmacogenomics, Faculty of Medicine, University of Maribor, Maribor, Slovenia

⁶Clinic of Pediatrics, University Medical Centre Maribor, Maribor, Slovenia

⁷COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital, University of Copenhagen, Copenhagen, Denmark

⁸Departments of Medicine, Bioengineering and Therapeutic Sciences, University of California, San Francisco, CA, USA

⁹Behavioral Sciences Institute, University of Puerto Rico, Medical Sciences Campus, San Juan, Puerto Rico

¹⁰Division of Pulmonary Medicine, Allergy, and Immunology, Children's Hospital of Pittsburgh of the University of Pittsburgh Medical Center, University of Pittsburgh, Pittsburgh, PA, USA

¹¹Department of Biological Sciences, National University of Singapore, Singapore, Singapore

¹²The Allergy & Immunology Division, Department of Paediatric Medicine, KK Children's Hospital, Singapore, Singapore

¹³Asthma Center, Connecticut Children's Medical Center, University of Connecticut Health Center, Hartford, CT, USA

¹⁴Department of Women's and Children's Health, University of Liverpool, Liverpool, UK

¹⁵Alder Hey Children's Hospital, Liverpool, UK

¹⁶Department of Pediatric Pneumology and Allergy, University Children's Hospital Regensburg (KUNO), Regensburg, Germany

¹⁷Department of Medical Informatics, Erasmus University Medical Center, Rotterdam, The Netherlands

¹⁸Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

¹⁹Department of Pediatrics, Sachs' Children's Hospital, Stockholm, Sweden

²⁰Academic Department of Paediatrics, Brighton and Sussex Medical School, Royal Alexandra Children's Hospital, Brighton, UK

²¹Division of Molecular & Clinical Medicine, School of Medicine, University of Dundee, Dundee, UK

²²CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain

²³Department of Molecular and Clinical Pharmacology, Institute of Translational Medicine, University of Liverpool, Liverpool, UK

²⁴Faculty for Chemistry and Chemical Engineering, University of Maribor, Maribor, Slovenia

²⁵Department of Human Genetics, Hannover Medical School, Hannover, Germany

²⁶Institute of Child Health, University College London, London, UK

²⁷Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden

²⁸Department of Biosciences and Nutrition, Karolinska Institutet, Stockholm, Sweden

²⁹The Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

³⁰Division of Pulmonary and Critical Care Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

³¹Division of Respiratory Medicine, Department of Pediatrics, Sainte-Justine University Hospital Center and University of Montreal, Montreal, QC, Canada

³²Child Health, University of Aberdeen, Aberdeen, UK
 Email: a.h.maitland@amc.uva.nl

REFERENCES

1. Chung KF, Godard P, Adelroth E, et al. Difficult/therapy-resistant asthma: the need for an integrated approach to define clinical phenotypes, evaluate risk factors, understand pathophysiology and find novel therapies. ERS Task Force on Difficult/Therapy-Resistant Asthma. *European Respiratory J*. 1999;13:1198-1208.
2. Moffatt MF, Gut IG, Demenais F, et al. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med*. 2010;363:1211-1221.
3. Tavendale R, Macgregor DF, Mukhopadhyay S, Palmer CNA. A polymorphism controlling ORMDL3 expression is associated with asthma that is poorly controlled by current medications. *J Allergy Clin Immunol*. 2008;121:860-863.
4. Berce V, Kozmus CEP, Potočnik U. Association among ORMDL3 gene expression, 17q21 polymorphism and response to treatment with inhaled corticosteroids in children with asthma. *Pharmacogenomics J*. 2013;13:523-529.
5. Farzan N, Vijverberg SJ, Andiappan AK, et al. Rationale and design of the multiethnic pharmacogenomics in childhood asthma consortium. *Pharmacogenomics*. 2017;18:931-943.
6. Leung TF, Ko FWS, Sy HY, Tsui SKW, Wong GWK. Differences in asthma genetics between Chinese and other populations. *J Allergy Clin Immunol*. 2014;133:42-48.

7. Toncheva AA, Potaczek DP, Schedel M, et al. Childhood asthma is associated with mutations and gene expression differences of ORMDL genes that can interact. *Allergy*. 2015;70:1288-1299.
8. Schmiedel BJ, Seumois G, Samaniego-Castruita D, et al. 17q21 asthma-risk variants switch CTCF binding and regulate IL-2 production by T cells. *Nat Commun*. 2016;7:13426.
9. Bisgaard H, Bønnelykke K, Sleiman PMA, et al. Chromosome 17q21 gene variants are associated with asthma and exacerbations but not

atopy in early childhood. *Am J Respir Crit Care Med*. 2009;179:179-185.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.